


RESEARCH ARTICLE

Seroprevalence of SARS-CoV-2 antibodies in Bali Province: Indonesia shows underdetection of COVID-19 cases by routine surveillance

Anak A. S. Sawitri^{1*}, Putu C. D. Yuliyatni¹, Putu A. S. Astuti¹, Emita Ajis^{2,3}, Endang B. Prasetyowati^{2,3}, Husni⁴, Juliette Morgan⁵, Jennifer Mika⁵, Catharina Y. Praptiningsih⁵, Amalya Mangiri⁵, Ester Mulyadi⁵, Rintis Noviyanti⁶, Leily Trianty⁶, William A. Hawley⁵

1 Department of Public Health and Preventive Medicine, Faculty of Medicine, Universitas Udayana, Denpasar, Bali, Indonesia, **2** Directorate of Health Surveillance and Quarantine, Ministry of Health Republic Indonesia, Jakarta Indonesia, **3** Gedung Adhyatma Kementerian Kesehatan Republik Indonesia, Jakarta, Indonesia, **4** Indonesia Field Epidemiology Secretariate, Jakarta Pusat, Indonesia, **5** US Centers for Disease Control and Prevention, Jakarta, Indonesia, **6** Eijkman Institute for Molecular Biology, Kota Jakarta Pusat, Daerah Khusus Ibukota Jakarta, Indonesia

* sagungawitri@gmail.com, sagung_sawitri@unud.ac.id



OPEN ACCESS

Citation: Sawitri AAS, Yuliyatni PCD, Astuti PAS, Ajis E, Prasetyowati EB, Husni, et al. (2022) Seroprevalence of SARS-CoV-2 antibodies in Bali Province: Indonesia shows underdetection of COVID-19 cases by routine surveillance. *PLOS Glob Public Health* 2(8): e0000727. <https://doi.org/10.1371/journal.pgph.0000727>

Editor: Gathsaurie Neelika Malavige, University of Sri Jayewardenepura, SRI LANKA

Received: February 25, 2022

Accepted: August 8, 2022

Published: August 31, 2022

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All data underlying the findings reported in this manuscript have been provided in [supplementary file](#) as part of the submitted article.

Funding: This work was supported by the U.S. Centers for Disease Control and Prevention [Cooperative Agreement Grant Number 6 NU2GGH001690-03-04 and 6 U18CK000443-05-01 to AASS]. The staff of CDC for Indonesia has

Abstract

The international tourist destination of Bali reported its first case of Coronavirus Disease 2019 or COVID-19 in March 2020. To better understand the extent of exposure of Bali's 4.3 million inhabitants to the COVID-19 virus, we performed two repeated cross-sectional serosurveys stratified by urban and rural areas. We used a highly specific multiplex assay that detects antibodies to three different viral antigens. We also assessed demographic and social risk factors and history of symptoms. Our results show that the virus was widespread in Bali by late 2020, with 16.73% (95% CI 12.22–21.12) of the population having been infected by that time. We saw no differences in seroprevalence between urban and rural areas, possibly due to extensive population mixing, and similar levels of seroprevalence by gender and among age groups, except for lower seroprevalence in the very young. We observed no difference in seroprevalence between our two closely spaced surveys. Individuals reporting symptoms in the past six months were about twice as likely to be seropositive as those not reporting symptoms. Based upon official statistics for laboratory diagnosed cases for the six months prior to the survey, we estimate that for every reported case an additional 52 cases, at least, were undetected. Our results support the hypothesis that by late 2020 the virus was widespread in Bali, but largely undetected by surveillance.

Introduction

Coronavirus Disease 2019 or COVID-19 is an infectious disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) [1]. Indonesia reported its first COVID-19 case

role on the development of study design, data collection, and preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

on 2 March 2020 in Jakarta [2, 3] with the international tourist destination of Bali reporting its first case shortly thereafter [4]. Reported COVID-19 cases in Indonesia generally increased throughout 2020 with some fluctuations [5]; similar trends were observed in Bali Province [6].

Published models from February 2020 suggested that COVID-19 reported cases in Indonesia were artificially low due to insufficient surveillance [7, 8] and imperfect reporting associated with a surveillance information system [9] that sometimes failed to report detected COVID-19 cases to provincial authorities. Early in the pandemic, Indonesian epidemiologists noted the need to increase testing for better tracking and control of the virus [10]. In May 2020, testing in Indonesia was limited, with only 225 tests administered per 1 million people, or 5.6% of the testing needed based upon projections at the time [10]. The problem of low testing was further compounded by inadequate tracking of confirmed cases, whereby data from the Bali Province Health Office showed that the average number of contacts traced per case was only nine persons in 2020, declining to less than 5 persons per case in October 2021 [11]. Thus, the true extent of virus spread in Bali was poorly understood.

Underreporting is exacerbated by the fact that most of Bali's reported cases were from hospitals treating symptomatic patients, which represent only a fraction of the clinical spectrum. The importance of pre-symptomatic COVID-19 transmission is well-documented whereby 44% of secondary cases were contracted before onset of COVID-19 symptoms [12]. The distinction between asymptomatic and pre-symptomatic transmission is problematic, whereby 'true' asymptomatic patients may be early in the process of developing symptoms [13]. A systematic review and meta-analysis revealed that asymptomatic persons have 50% lower probability to transmit SARS-CoV-2 compared to those with symptoms [14].

In addition to assessing the extent of exposure, seroprevalence surveys are useful for monitoring trends, geographic distribution, and targeting interventions [15]. At the time of this survey, similar seroprevalence studies had been reported elsewhere [16, 17] but none had been conducted in Bali. Further, our repeated cross-section study design has the potential to assess the rapidity of spread of the virus. Our expectation is that our results will inform public health measures for better surveillance and control [15].

Material and methods

Ethics and permission

Ethical approval was received from the Ethics Committee of Faculty of Medicine Udayana University on 20 July 2020 prior to the first round of the survey, and an amendment was proposed for measurement of blood pressure and blood sugar for adults and determining blood type for children as incentives for round 2; approval for the amendment was granted on 7 November 2020. Written permission from the Governor of Bali Province had been granted before the data collection. All survey procedures and related information were explained to eligible subject or their legal representative, by enumerators on behalf of principal investigator prior the data collection. We interviewed and took blood samples only from parents and eligible respondents who provide their written consent.

Setting and study design

Bali Province has a population of 4.3 million with 785,000 people in its capital, Denpasar [18]. Population mobility is high in and out of Bali Province, between districts, and between urban and rural areas [19]. For this survey, Denpasar, with a population density of 6,360 people/km² was considered an urban area, while the remainder of the island with a population density around or below of 7.50 people/km² was considered rural [20].

Cross-sectional sero-surveys were conducted in two rounds between 14 October– 24 December 2020. Specimen collection for round 1 occurred from 14 October– 14 November and for round 2 from 15 November– 24 December. Urban (Denpasar) compared to rural (outside Denpasar) stratification was done. Multi-stage cluster random sampling of census enumeration areas was performed.

Population and sample

Sample size and power. The survey was designed to provide an estimate of seroprevalence of SARS-CoV-2 in Bali Province, in urban Denpasar, and in rural areas outside of the capital. The minimum required sample size was calculated using this formula.

$$\text{Minimum required sample size} = DE \times \frac{Z_{1-\alpha/2}^2 \times P \times (1 - P)}{d^2}$$

We assumed that the expected prevalence of antibodies to SARS-CoV-2 was higher in Denpasar (7%) than in the rest of the island (5%). We also assumed a conservative design effect (DE) of 2 to account for an expected increase of variance due to clustering effect, a response rate of 75%, and a desired confidence level of estimate $1-\alpha/2 = 95\%$ ($Z_{1-\alpha/2} = 1.96$) and a 2% margin of error (d), resulting in a sample size of 1,555 in urban Denpasar and 1,138 in rural areas (total of 2,693 individuals) per survey. With an average expected number of persons per household of 3.46 in Denpasar and of 3.89 in the non-Denpasar areas [21], the number of households expected to be visited was 450 in urban and 293 in rural areas (total of 743 households) per survey.

Sample, household (HH) and participant selection. A household was defined as a group of persons who reside in the same place and prepare meals together. All members of selected households who were ≥ 1 year of age were eligible if informed consent was obtained by either the participant of the parent or guardian.

The National Statistics Office provided enumeration areas (census blocks based upon a March 2020 national social and economic survey) for Bali as the sampling frame. Twenty HHs were selected per census block, with 23 from Denpasar and 15 from rural districts, for a total of 38 blocks.

A two-stage systematic random sampling strategy was applied. In stage 1, census blocks were selected in urban and rural areas, while in stage two, 20 HHs in each census block were selected after performing stratification by educational level of the head of HH. The samples were mutually independent between surveys, whereby the census blocks with odd serial numbers were used as the sampling frame for the round 1 survey, while census blocks with even serial numbers were used for round 2. A sample reserve of 20% or 4 HHs per census block was prepared. Selection of backup samples was carried out by systematic sampling from the list of HHs not selected in the original procedure.

Data collection

Training. A technical guide for fieldwork including all survey procedures was prepared and reviewed during two days of training for the eight survey teams. The survey teams were monitored by project staff throughout the survey.

Questionnaire. A structured questionnaire in the Indonesian language was developed based on previous behavioral and risk factor assessments for SAR-CoV2 exposure [22–24]. Basic demographic and economic information (number of HH members, HH income, age, gender, education, and marital status) was obtained along with a history of symptoms during the last 6 months. We also developed a non-response form for refusers. The interview

questions were pilot tested and evaluated by local experts to ensure that they were culturally appropriate and could be understood by a layperson with a primary education. The questionnaire was revised after piloting in enumeration areas not selected for the survey.

Interview. Survey teams visited HHs in coordination with local leaders and guides from the district statistical office. Interview responses were recorded on a password protected hand-held device that uploaded to a secure database upon completion of the questionnaire. Paper format was used in areas where internet access was limited.

Blood sampling and processing. Dried blood spot (DBS) samples were collected by finger prick and spotted onto Whatman 903 filter paper. An identification number was placed on each filter paper. Each participant's data included a unique identifier (barcoded label). Data collection in each census block was on average completed in four days.

DBS specimens were registered in an electronic log book for tracking. Blood spots were dried overnight at room temperature until uniformly dark brown with no red color visible. The DBS were then stored and shipped at 4–8°C via air to the Eijkman Institute for Molecular Biology in Jakarta, where they were stored at 20°C.

SARS-CoV-2 antibody testing was performed on a Luminex MAGPIX instrument using the Tetracore FlexImmArray SARS-CoV-2 Human IgG Antibody assay [25]. This test detects antibodies to three different antigenic sites on the virus: spike (S), nucleoprotein (NP) and hybrid using a Multiplex Bead technology. The test uses 7 microspheres, with three detecting antibodies to different SARS-CoV-2 antigens, while 4 are internal controls. Every 96-well test plate was used for testing up to 90 samples in one test run. Each test run included SARS-CoV-2 negative control serum, positive control serum, and calibrator in duplicates. Antibody response was determined qualitatively using the ratio of mean fluorescence intensity (MFI) target antigen/MFI calibrator. Results were considered SARS-CoV2 human IgG positive if all three target antigens had a ratio of ≥ 1.2 , negative if the ratio was ≤ 0.9 , and indeterminate if between this range. Specimens with indeterminate results were re-tested; if the second test was indeterminate the specimen was classified as negative. A positive result implied past exposure to the virus.

Data analysis

Weighting was performed by the National Statistical Office to account for sampling design, response rate, and stratification. Descriptive analysis was conducted for characteristics of the HHs and respondents for round 1, round 2 and combined rounds. The overall prevalence of SARS-CoV-2 antibodies was calculated using weighted and unweighted data, per round and for both rounds combined. To estimate a population value, we calculated a range of estimates as a percentage, whereby a population mean lies between an upper and lower interval. Prevalence was calculated for rural and urban areas, by socio-demographic parameters, and by presence or absence of reported symptoms. Sensitivity analysis was not carried out, as an unpublished report showed that the Tetracore test kit had a sensitivity of 89% and specificity of 100% [25]. Raw data used for analysis is provided in (S1 Data).

In Bali province, monitoring of COVID-19 cases is performed through active and passive surveillance systems. The passive system records the number of confirmed infections, deaths, and recovered hospitalized COVID-19 people from public health centers, local and central public hospitals, and some private hospitals in Bali Province. Meanwhile, active surveillance records cases found from contact tracing and screening at the entry points such as the airport and harbour. Information from active and passive surveillance is entered to the single sign on (SSO) system. The system was developed in mid June 2020 and was used until September 2021. Since then, COVID-19 data was transferred to a central information system called 'New

All Record' (NAR). The New All Record system was established in April 2021 [26]. The possibility of a gap between detected and reported COVID-19 persons in this surveillance system is beyond the scope of our study.

Using the SSO data from 1 June—30 November 2020 (<https://infocorona.baliprov.go.id/>) for which period we assumed that infections would produce detectable antibodies during the time of our survey, we compared the estimated seroprevalence of SARS-CoV-2 with reported cumulative COVID-19 cases in Bali Province and calculated number of missed infections during this period.

Results

Table 1 presents the characteristics of census blocks being surveyed. Among 38 targeted census blocks, one census block was not surveyed during round 1 due to refusal by local authorities; in round 2, all 38 census blocks were successfully surveyed. The acceptance rate of HHs and respondents was slightly higher in rural than in urban areas, and was higher in round 2 than in round 1. The acceptance rate for DBS collection was low in both rounds, but was higher in rural areas and in round 2. Very few samples failed quality control for the lab assays, with only 1.8% failing in round 1 and 1.1% failing in round 2. Samples that did not pass QC were excluded from the analysis.

Table 2 presents the sociodemographic and household characteristics of respondents consenting to interview and blood sampling. Respondents in the 15–54 year old age group were more likely to consent to interview than those younger or older than this group. Gender and marital status were not associated with consent to interview. The median level of education was through high school, with mean household monthly income in both surveys about 3 million IDR (215 USD). While most households have ventilation, only 20% were airconditioned. Blood samples were obtained from a low proportion of children (1–4 y.o., and 5–14 y.o.) in both rounds, with an increase in round 2.

Table 3 shows details of seroprevalence for each survey round. The combined overall prevalence for both surveys (N = 2,545), without and with weighting were 17.5 (95% CI 16.01–18.96) and 16.73 (95% CI 12.22–21.12), respectively.

Contrary to expectation, neither survey round showed a statistically significant difference in prevalence between urban versus rural areas. Prevalence based on socio-demographics also

Table 1. Characteristics of survey census blocks.

Districts	Census Block type			Description of Household			Respondent characteristics				Average number of respondents per HH	
	Planned		Surveyed	Approached		Surveyed	Eligible HH members		Number interviewed			Blood Samples obtained
	N	N	%	N	N	%	N	N	%	N		%
Round 1												
Bali	38	37	97.4	793	742	93.6	2,392	2,353	98.37	1,175	49.94	3.22
Urban	23	23	100	504	463	91.8	1,396	1,362	97.56	591	43.39	3.01
Rural	15	14*	93.3	289	279	96.5	996	991	99.50	584	58.93	3.57
Round 2												
Bali	38	38	100	785	760	96.8	2,089	2,076	99.38	1,407	67.77	2.75
Urban	23	23	100	484	460	95.0	1,222	1,211	99.00	809	66.80	2.66
Rural	15	15	100	301	300	99.7	867	865	99.77	645	74.57	2.89

Note: Urban = Denpasar, Rural = 8 districts non-Denpasar; eligible member of HH: number of family member \geq 1 years old plus non family member who stay in the same house, average respondents per HH: eligible member of HH/ selected HH Surveyed.

<https://doi.org/10.1371/journal.pgph.0000727.t001>

Table 2. Characteristics of survey participants.

Socio-demographic characteristics	Interviewed		Blood Sample obtained		Proportion of those interviewed with blood sample obtained	
	Round 1 (N = 2,351)	Round 2 (N = 2076)	Round 1 (N = 1,175)	Round 2 (N = 1,407)	Round 1	Round 2
	Freq (%)	Freq (%)	Freq (%)	Freq (%)	%	%
Age Group						
1–4	126 (5.36)	86 (4.14)	6 (0.51)	18 (1.28)	4.76	20.93
5–14	335 (14.25)	275 (13.25)	33 (2.81)	45 (3.20)	9.85	16.36
15–24	458 (19.48)	326(15.70)	63 (5.36)	87(6.18)	13.76	26.69
25–34	332 (14.12)	290(13.97)	98 (8.34)	108 (7.68)	29.52	37.24
35–44	366(15.44)	343 (16.52)	149 (12.68)	112 (7.96)	40.82	32.65
45–54	363 (15.44)	390(18.79)	85 (7.23)	96 (6.82)	23.42	24.62
55–64	223 (9.49)	207(9.97)	88 (7.49)	108 (7.68)	39.46	52.17
65+	148 (6.30)	159 (7.66)	97 (8.26)	114 (8.10)	65.54	71.70
Sex						
Male	1.171 (49.77)	1.033 (49.76)	572 (48.68)	651 (46.27)	48.85	63.02
Female	1.182 (50.23)	1.043 (50.24)	603 (51.32)	756 (53.73)	51.02	72.48
Marital status						
Not married	1,000 (42.50)	756 (36.42)	380 (32.34)	426 (30.28)	38.00	56.35
Married	1,310 (55.67)	1,262 (60.79)	772 (65.70)	929 (66.03)	58.93	73.61
Divorce	43 (1.83)	58 (2.79)	23 (1.96)	53 (3.70)	53.49	91.38
Education						
Yet/never attended school	265 (11.26)	181 (8.72)	73 (6.21)	91 (6.47)	27.55	50.28
Not completed ES	153 (6.50)	156 (7.51)	71 (6.04)	82 (5.83)	46.41	52.56
Elementary school (ES)	453 (19.25)	519 (25.00)	230 (19.57)	364(25.87)	50.77	70.13
Junior high school	345 (14.66)	298 (14.35)	202 (17.19)	226 (16.06)	58.55	75.84
High school	783 (33.28)	656 (31.60)	439 (37.36)	461(32.76)	59.90	70.27
Diploma	94 (3.99)	81 (3.90)	43 (3.66)	55 (3.91)	45.74	67.90
Bachelor (D4/S1)	240 (10.20)	174 (8.38)	106 (9.02)	123 (8.74)	44.17	70.69
Master/Doctor	20 (0.85)	11 (0.53)	11 (0.94)	5 (0.35)	55.00	45.45
HOUSEHOLD*	N = 740	N = 760				
Household income (million IDR)						
Mean (SD)	3,061 (4.18)	3.135 (6.28)				
Range (million/month)	0–54	0–88				
Ventilation in house	N = 740	N = 760				
Available	678 (91.62)	678 (89.21)				
Not available	62 (8.32)	82 (10.79)				
Air conditioning in house	N = 742	N = 760				
Available	166 (22.37)	147 (19.34)				
Not available	576 (77.63)	613 (80.66)				

*Characteristics of participating survey households

<https://doi.org/10.1371/journal.pgph.0000727.t002>

showed no significant difference between sex, age group, education level, marital and employment status in either survey round. Respondents with symptoms were more likely to be seropositive than asymptomatic individuals.

As shown in “Fig 1”, seroprevalence results were extrapolated to estimate the total number of infections in Bali compared to the actual number of infections reported by the surveillance system of the Government of Bali. The results show that only about 2% of infections were reported, with 52 cases likely occurring for each laboratory confirmed reported case.

Table 3. Seroprevalence of SARS-CoV-2.

Variable	Round 1 (N = 1,154)				Round 2 (N = 1,391)			
	N total	%	95% CI		N total	%	95% CI	
			Lower	Upper			Lower	Upper
Overall Prevalence								
without weighting		16.72	14.57	18.88		18.12	16.09	20.14
with weighting		18.04	9.98	26.10		15.71	11.15	20.28
Area								
Urban	583	13.08	8.61	17.55	773	17.67	12.16	23.17
Rural	571	19.13	8.33	29.92	618	15.26	9.25	21.26
Sex								
Male	562	17.79	9.02	26.55	648	15.09	8.66	21.52
Female	592	18.29	9.94	26.63	743	16.28	11.50	21.06
Age group (year)								
1–4	7	0	-	-	18	9.4	0.00	28.29
5–14	92	25.89	12.15	39.63	132	14.99	5.84	24.13
15–24	246	15.69	6.16	25.22	220	16.95	9.24	24.67
25–34	167	17.98	12.06	23.89	202	16.68	8.19	25.18
35–44	201	16.67	7.34	26.02	249	14.11	7.68	20.55
45–54	213	18.76	7.56	29.96	290	18.10	8.61	27.60
55–64	137	18.62	5.39	31.85	152	16.74	7.57	5.91
65+	91	18.39	2.11	34.69	1128	11.81	4.47	19.15
Education level								
Illiterate/not complete ES	142	20.58	8.87	32.28	167	15.55	4.59	26.51
ES–High School	854	17.02	10.03	24.00	1,041	16.59	10.97	22.20
Diploma/University	158	20.88	9.84	40.85	183	9.66	2.52	16.81
Marital status					N = 1,350*			
Not married	374	18.14	10.45	25.82	421	15.06	8.91	21.21
Married	758	18.40	9.916	27.65	918	16.03	10.13	21.95
Divorced	22	5.06	0.02	12.11	11	16.25	5.68	26.82
Working status	N = 1,153*							
Unemployed	522	18.02	8.89	27.23	625	13.39	8.64	18.15
Employed	631	18.06	10.00	26.03	766	17.84	11.43	24.24
Symptom status	N = 1,140*				N = 1,384*			
No symptom	919	16.20	9.86	22.53	1,091	13.72	8.99	18.45
Common symptoms (fever, cough, rhinitis, dyspnea)	100	23.23	6.54	39.91	117	20.65	7.72	33.57
Less common symptoms (others)	75	27.30	1.21	53.40	135	20.00	7.58	32.44
With symptoms (less/common)	46	34.76	7.00	62.50	41	26.45	8.35	44.55

Note: Seroprevalence of SARS-CoV-2 based on area, gender, sex, marital, education and symptoms are weighted; * contain missing data

<https://doi.org/10.1371/journal.pgph.0000727.t003>

Nearly three fourths of respondents reported no symptoms of SARS-CoV-2 infection in the last six months, as shown in Table 4. The most common symptoms reported were fever, headache, and rhinitis.

Discussion

This article reports results of a SARS-CoV-2 repeated cross-sectional serosurvey conducted in Bali Province in late 2020 [27]. The study was conducted relatively early in the pandemic, yet a high prevalence of SARS-CoV-2 antibodies was detected with a round 1 estimate of 18.04%




Fig 1. Comparison of reported case and estimated cases of SARS-CoV-2 infection based on serosurvey results.

<https://doi.org/10.1371/journal.pgph.0000727.g001>

(95% CI 9.98–26.10), a round 2 estimate of 15.71% (95% 11.25–20.28), and an overall combined prevalence of 16.73% (95% CI 12.22–21.12).

The results of this survey are similar to those obtained in July 2020 in India [16, 28], lower than results from a survey in Iran in April 2020 [29], but much higher than results from a systematic review for Asia and Southeast Asia for January–December 2020 showing seroprevalence of only 0.6% (0.3–1.4%) [27]. Within Indonesia, results from this survey were significantly lower than those obtained in Jakarta in March 2021, showing seroprevalence of 44.6% using the same laboratory methods [30]. However, the prevalence reported in the Bali survey was higher than that reported in during a comparable time in East Java (11%) [31].

Table 4. Symptoms reported by individuals positive for SARS-CoV-2 antibody within the last six months.

Symptoms	Round 1 (n = 192)		Round 2 (n = 250)		All Round (n = 442)	
	n	%	n	%	n	%
With symptoms						
No	140	72.9	180	72.0	320	72.4
Yes	52	27.1	70	28.0	122	27.6
Types of symptoms						
Fever	20	10.4	20	8.0	40	9.0
Rhinitis	10	5.2	13	5.2	23	5.2
Cough	10	5.2	11	4.4	21	4.8
Dyspnea	4	2.1	3	1.2	7	1.6
Sub-fever	3	1.6	13	5.2	16	3.6
Headache	13	6.8	21	8.4	34	7.7
Myalgia	7	3.6	4	1.6	11	2.5
Athralgia	11	5.7	4	1.6	15	3.4
Fatigue	11	5.7	2	0.8	13	2.9
Nausea	8	4.2	3	1.2	11	2.5
Vomiting	3	1.6	2	0.8	5	1.1
Diarrhea	3	1.6	5	2.0	8	1.8
Lose of appetite	4	2.1	1	1	5	1.1

<https://doi.org/10.1371/journal.pgph.0000727.t004>

The estimated level of SARS-CoV-2 prevalence from this survey implies that the surveillance system in Bali detected and reported an extremely low proportion of positive cases as shown in “Fig 1”. Meanwhile in Jakarta, a more massive testing effort was able to detect and report a higher proportion (8.1%) of infections [30]; however, this is still far below the required number by WHO [10, 11]. This condition occurred not only in Indonesia but elsewhere, as one systematic review has reported that numbers of infections estimated by serosurveys are always higher than actual number of cases detected [27]. The low proportion of positive cases detected by the surveillance system reflects the fact that neither Bali’s testing rate nor contact tracing efforts meet WHO standards [10, 11]. The weakness of surveillance systems to detect COVID-19 has also been reported in Africa [32]. Suboptimal reporting and monitoring of COVID-19 cases may create a false impression of decreasing of COVID-19 incidence [32]. Critical assessment of COVID-19 data in Indonesia [33] showed that the surveillance system only reported the confirmed, recovered and fatal cases and did not report suspected cases who died. It also missed the geographic and demographic details at the national, provincial and district level, as has been reported in Africa [32]. Suboptimal surveillance may mis-direct policy-making and control strategies [32] and may diminish the effectiveness of policy initiatives at the local level [33] and in the worst case adversely affect morbidity and mortality rates.

This survey was conducted using a standard methodology. Overestimation of prevalence due to clustering of cases within families was low, given that only 7% of families had more than one positive individual. As the laboratory test used requires positivity for antibodies to three different antigens, its specificity is high (100%) while sensitivity is 89% [15], leading to possible underestimation of seroprevalence.

Symptoms which were possibly related to COVID-19 were reported in 34.76% (95%CI 7.00–62.50) of respondents in round 1 and 26.45% (95% CI 8.35–44.45) of respondents in round 2. In contrast, seroprevalence in those not reporting symptoms was about half this level, though the differences were not statistically significant. In Iran, the proportion of seropositive asymptomatic individuals was much higher (57.2%) [17]. Even if the potential spread from asymptomatic cases is low [13], their relatively high numbers leads to significant transmission risk [34, 35], particularly in the context of weak active case detection and contract tracing.

Results from this study showed that 72% of seropositive respondents reported no symptoms whatsoever. Those numbers were probably too high [13] as a recent meta-analysis estimated the proportion of asymptomatic cases at 17% (95%CI 14–20%) [14], though several other studies showed wider variation [17, 36]. This study is subject to recall bias, as we asked about symptoms occurring in the past six months, whereby mild illness may have been forgotten or judged as not being ill at all. Further, at the time of the interview and DBS collection, some participants may have been subclinical or pre-symptomatic [14], which have shown high risk of transmission [13, 36]. Therefore, results from this survey support the conclusion that by focusing on symptomatic individuals, programs lose the opportunity to prevent transmission from asymptomatic individuals [13, 14].

No seroprevalence differences was found between the two survey rounds. During the first round of data collection, the response rate for blood draw was very low for various reasons (e.g. fear of diagnosis with COVID-19, fear of needles, too young for blood draw etc.). We attempted to increase the response rate by re-socialization to the mayor, head of districts and sub-districts, head of villages and sub-villages; and provided additional incentives, such as blood type test (for children), blood glucose test and blood pressure test (for adult respondents) for their participation. These measures helped to to increase the acceptance for blood draw in the second survey, resulting in a narrowing of the 95% confidence interval of SARS-CoV-2 seroprevalence in the second compare to first round survey. Nevertheless, analysis of sociodemographic characteristics of respondents for the two rounds showed that acceptance

of blood draw for females was significantly higher than male ($p < 0.00$), and for rural areas was significantly higher than for urban areas ($p < 0.00$). Related to symptoms and acceptance for blood draw, the percentage of those reporting headache and dyspnea were higher in those providing blood samples compared those not reporting these symptoms. However, for other symptoms, we found no differences among those with and without symptoms, making us confident that differences in blood collection success between survey rounds did not result in significant bias.

We found no difference in prevalence between urban Denpasar and the rest of rural Bali. This may be because road transport and access throughout Bali is relatively good [13]. In addition, movement between rural and urban areas is common, as most respondents (round 1 = 71.9% and round 2 = 69.7%) reported participating in traditional ceremonies in their home villages while working or residing elsewhere.

This study has some weaknesses. First, response rate was low for DBS collection, with responses particularly low among very young participants. Second, as symptoms were reported by recall, the results are subject to bias. Nonetheless, results do show that the virus had spread substantially in Bali by late 2020, in contrast to findings from official statistics.

Supporting information

S1 Data. Data of seroprevalence of SARS-CoV-2 in Bali.
(XLS)

Acknowledgments

We thank our study coordinators and enumerators who worked hard to collect data in the field, as well as all respondents for their participation. We also thank to Indonesia Statistic Office for providing sampling frame and thank the Government of Bali for permission to publish these data.

Author Contributions

Conceptualization: Anak A. S. Sawitri, Putu C. D. Yuliyatni, Juliette Morgan, Catharina Y. Praptiningsih, Amalya Mangiri, Ester Mulyadi, Rintis Noviyanti, Leily Trianty.

Data curation: Anak A. S. Sawitri, Putu C. D. Yuliyatni, Putu A. S. Astuti.

Formal analysis: Anak A. S. Sawitri, Putu C. D. Yuliyatni, Putu A. S. Astuti.

Funding acquisition: Juliette Morgan, Jennifer Mika, William A. Hawley.

Methodology: Anak A. S. Sawitri, Putu C. D. Yuliyatni, Jennifer Mika, Catharina Y. Praptiningsih, Amalya Mangiri, Ester Mulyadi, Rintis Noviyanti, William A. Hawley.

Project administration: Emita Ajis, Endang B. Prasetyowati, Husni.

Resources: Emita Ajis, Endang B. Prasetyowati, Husni, Juliette Morgan, Catharina Y. Praptiningsih, Amalya Mangiri, Ester Mulyadi, Rintis Noviyanti, Leily Trianty.

Supervision: Putu C. D. Yuliyatni, Emita Ajis, Endang B. Prasetyowati, Husni, Jennifer Mika.

Validation: Putu A. S. Astuti, Rintis Noviyanti, Leily Trianty, William A. Hawley.

Visualization: Putu A. S. Astuti.

Writing – original draft: Anak A. S. Sawitri.

Writing – review & editing: Anak A. S. Sawitri, Putu C. D. Yuliyatni, Putu A. S. Astuti, Emita Ajis, Endang B. Prasetyowati, Juliette Morgan, Jennifer Mika, Catharina Y. Praptiningsih, Amalya Mangiri, Ester Mulyadi, Leily Trianty, William A. Hawley.

References

1. Ludwig S, Zarbock A. Coronaviruses and SARS-CoV-2: A Brief Overview. *Anesth Analg* [Internet]. 2020; 131(1):93–6. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7173023/> <https://doi.org/10.1213/ANE.0000000000004845> PMID: 32243297
2. World Health Organization. Media Statement on confirmed COVID-19 cases. *World Heal Organ* [Internet]. 2020;2020–2. Available from: <https://www.who.int/indonesia/news/detail/02-03-2020-media-statement-on-covid-19>
3. World Health Organization Indonesia. Corona Virus Disease 2019 Situation Report [Internet]. Vol. Maret, Situation Report. 2020. p. 1–8. Available from: https://www.who.int/docs/default-source/search/indonesia/covid19/who-indonesia-situation-report-1.pdf?sfvrsn=6be5b359_0
4. Presiden RI. Keppres No. 12 Tahun 2020 Tentang Penetapan Bencana Nonalam Penyebaran Corona Virus Disease 2019 Sebagai Bencana Nasional. 2020 p. 1–2.
5. John Hopkins University. CSSEGISandData [Internet]. 2020 [cited 2021 Oct 1]. Available from: <https://github.com/CSSEGISandData/COVID-19>
6. Indonesian Covid-19 Control Task Force (Satuan Tugas Penanggulangan Covid-19 Indonesia). Peta Sebaran (Distribution Map) [Internet]. National Government of Indonesia. 2020 [cited 2021 Oct 1]. Available from: <https://covid19.go.id/peta-sebaran>
7. Salazar PM De, RN, A T, C B, Lipsitch M. Using predicted imports of 2019-nCoV cases to determine locations that may not be identifying all imported cases. *medRxiv*. 2020;(165):1–13.
8. Syakriah A. Testing gap across provinces slows down COVID-19 response in Indonesia. *The Jakarta Post* [Internet]. 2020 Jul 22; Available from: <https://www.thejakartapost.com/news/2020/07/22/testing-disparity-among-regions-a-challenge-for-covid-19-response.html>
9. Barsasella D, Iman AT, Fadly F, Uddin M, Mohammed A, Shaik T, et al. Information Flow and Data Gaps in COVID-19 Recording and Reporting at National and Provincial Levels in Indonesia. *Healthcare* [Internet]. 2022 Jan 20; 10(2):204. Available from: <https://www.mdpi.com/2227-9032/10/2/204> <https://doi.org/10.3390/healthcare10020204> PMID: 35206818
10. Ariawan I, Riono P, Farid MN, Jusril H, Wahyuningsih W, Ali PB, et al. Proyeksi COVID-19 di Indonesia. 2021. 52 p.
11. Bali Province Health Office. Perkembangan Covid-19 Bali (The Covid-19 Situation Progress in Bali Province) [Internet]. 2020 [cited 2021 Sep 25]. Available from: <https://www.diskes.baliprov.go.id/perkembangan-covid-19-bali/>
12. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med*. 2020; 26(5):672–5. <https://doi.org/10.1038/s41591-020-0869-5> PMID: 32296168
13. Pollock AM, Lancaster J. Asymptomatic transmission of covid-19. *BMJ*. 2020; 371:19–20.
14. Byambasuren O, Cardona M, Bell K, Clark J, McLaws ML, Glasziou P. Estimating the extent of asymptomatic COVID-19 and its potential for community transmission: Systematic review and meta-analysis. *J Assoc Med Microbiol Infect Dis Canada*. 2020; 5(4):223–34.
15. Peeling RW, Olliaro PL. The time to do serosurveys for COVID-19 is now. www.thelancet.com/respiratory [Internet]. 2020; 8(September):836–7. Available from: www.thelancet.com/respiratory
16. Selvaraju S, Kumar MS, Thangaraj JWV, Bhatnagar T, Saravanakumar V, Kumar CPG, et al. Population-based serosurvey for severe acute respiratory syndrome coronavirus 2 transmission, Chennai, India. *Emerg Infect Dis*. 2021; 27(2):586–9. <https://doi.org/10.3201/eid2702.203938> PMID: 33496222
17. Shakiba M, Nazemipour M, Heidarzadeh A, Mansournia MA. Prevalence of asymptomatic COVID-19 infection using a seroepidemiological survey. *Epidemiol Infect*. 2020;4–6. <https://doi.org/10.1017/S0950268820002745> PMID: 33183367
18. Badan Pusat Statistik Provinsi Bali. Berita Resmi Statistik No. 08/01/51/Th. II. WwwBaliBpsGold. 2021; (08):1–13.
19. Damuri, Yose Rizal, Prabaning Tyas, Lionel Priyadi, Stella Kusumawardhani EKY. Tracking the Ups and Downs in Indonesia's Economic Activity During COVID-19 Using Mobility Index: Evidence from Provinces in Java and Bali [Internet]. Jakarta; 2021. (ERIA Discussion Paper). Report No.: 385. Available from: <https://www.eria.org/uploads/media/discussion-papers/ERIA-Research-on-COVID-19/>

[Tracking-the-Ups-and-Downs-in-Indonesia's-Economic-Activity-During-COVID-19-Using-Mobility-Index.pdf](#)

20. National Statistic Office. Bali dalam Angka (Bali in Numbers). [Internet]. Denpasar; 2020. Available from: <https://bali.bps.go.id/publication/2020/04/27/c2d74be21e44651b07e2658e/provinsi-bali-dalam-angka-2020.html>
21. Statistics Indonesia. Rata-rata Banyaknya Anggota Rumah Tangga 2014–2016 (Average of House Hold Member 2014–2016) [Internet]. 2020 [cited 2020 Jul 12]. Available from: <https://www.bps.go.id/indicator/12/148/1/rata-rata-banyaknya-anggota-rumah-tangga.html>
22. Bi Q, Lessler J, Eckerle I, Lauer SA, Kaiser L, Vuilleumier N, et al. Household Transmission of SARS-CoV-2: Insights from a Population-based Serological Survey. medRxiv Prepr. 2020;(November 2020).
23. Bhagat RK, Davies Wykes MS, Dalziel SB, Linden PF. Effects of ventilation on the indoor spread of COVID-19. *J Fluid Mech* [Internet]. 2020 Nov 25; 903:F1. Available from: https://www.cambridge.org/core/product/identifier/S002211202000720X/type/journal_article <https://doi.org/10.1017/jfm.2020.720> PMID: 34191877
24. Mendoza-Jiménez M-J, Hannemann T-V, Atzendorf J. Behavioral Risk Factors and Adherence to Preventive Measures: Evidence From the Early Stages of the COVID-19 Pandemic. *Front Public Heal* [Internet]. 2021 Jun 9; 9. Available from: <https://www.frontiersin.org/articles/10.3389/fpubh.2021.674597/full> PMID: 34178929
25. Kaitlin F Mitchell, Christina M Carlson, Douglas Nace, Brian S Wakeman, Jan Drobeniuc, Glenn P Niemeyer, et al. Evaluation of a multiplex bead assay for detection of IgG antibodies to SARS-CoV-2 (draft). -. 2021; 23.
26. Indonesia Ministry of Health. Pemerintah integrasikan data kesehatan dengan aplikasi Pedulilindungi untuk mencegah pemalsuan hasil tes COVID-19 sebagai syarat perjalanan (Government integrate health data in to Pedulilindungi application to prevent fake COVID-19 test result as a require [Internet]. 2021 [cited 2022 Feb 23]. Available from: <https://sehatnegeriku.kemkes.go.id/baca/rilis-media/20210704/0738021/pemerintah-integrasikan-data-kesehatan-dengan-aplikasi-pedulilindungi-untuk-mencegah-pemalsuan-hasil-tes-covid-19-sebagai-syarat-perjalanan/>
27. Bobrovitz N, Arora RK, Cao C, Boucher E, Liu M, Donnici C, et al. Global seroprevalence of SARS-CoV-2 antibodies: A systematic review and metaanalysis. *PLoS One* [Internet]. 2021; 16(6 June):1–21. Available from: <http://dx.doi.org/10.1371/journal.pone.0252617>
28. Prakash O, Solanki B, Sheth JK, Joshi B, Kadam M, Vyas S, et al. Assessing seropositivity for IgG antibodies against SARS-CoV-2 in Ahmedabad city of India: A cross-sectional study. *BMJ Open*. 2021; 11(1):1–8.
29. Shakiba M, Nazemipour M, Salari A, Mehrabian F, Hashemi Nazari SS, Rezvani SM, et al. Seroprevalence of SARS-CoV-2 in Guilan Province, Iran, April 2020. *Emerg Infect Dis*. 2021; 27(2):636–8. <https://doi.org/10.3201/eid2702.201960> PMID: 33349310
30. Kesehatan D, Dki P, Fkm-ui TP, Eijkman L, Indonesia CDC. Prevalensi Antibodi Positif SARS CoV-2 di DKI Jakarta. 2021;
31. Megasari NLA, Utsumi T, Yamani LN, Juniastuti, Gunawan E, Furukawa K, et al. Seroepidemiological study of SARS-CoV-2 infection in East Java, Indonesia. Di Gennaro F, editor. *PLoS One* [Internet]. 2021 May 6; 16(5):e0251234. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0251234> PMID: 33956869
32. Aborode AT, Hasan MM, Jain S, Okereke M, Adedeji OJ, Karra-Aly A, et al. Impact of poor disease surveillance system on COVID-19 response in africa: Time to rethink and rebuilt. *Clin Epidemiol Glob Heal* [Internet]. 2021 Oct; 12:100841. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8330137/> <https://doi.org/10.1016/j.cegh.2021.100841> PMID: 34368503
33. Januraga PP, Harjana NPA. Improving Public Access to COVID-19 Pandemic Data in Indonesia for Better Public Health Response. *Front Public Heal* [Internet]. 2020 Nov 24; 8. Available from: <https://www.frontiersin.org/articles/10.3389/fpubh.2020.563150/full> PMID: 33330311
34. Zhang Jinzhong, Zhou Pu HD, Wenchao W, Cui C, Ran Z, Kexin X, et al. Investigation on a cluster of novel coronavirus pneumonia in a supermarket in Liaocheng. *Chinese J Epidemiol*. 2020; 41(12):2024–8.
35. Wirawan A, Januraga PP. Forecasting COVID-19 Transmission and Healthcare Capacity in Bali, Indonesia. *J Prev Med Public Heal*. 2020; 53(3):158–63. <https://doi.org/10.3961/jpmph.20.152> PMID: 32498137
36. Johansson MA, Quandelacy TM, Kada S, Prasad PV, Steele M, Brooks JT, et al. SARS-CoV-2 Transmission from People without COVID-19 Symptoms. *JAMA Netw Open*. 2021; 4(1):4–11. <https://doi.org/10.1001/jamanetworkopen.2020.35057> PMID: 33410879